

MODELING AN ACTIVE (!!) EXPLOSIVE VOLCANO

Molly F. Miller and Thomas C. Moyer
Geology Department
Vanderbilt University
Nashville, TN 37235

Level: Grades 3 - 6

Estimated time required: 30 minutes

Anticipated Learning Outcomes

- Students will explain how the build-up of gas from dissolving alka seltzer causes the lid of a film cannister to blow off.
- Students will explain that build-up of gas pressure causes eruption of explosive volcanoes, and that the pressure comes from heating of dissolved gases in the magma.
- Students will delineate the similarities and differences between the model and actual volcano.
- Students will verbalize excitement about learning about volcanoes.

Introduction

This activity is an active simulation of an explosive volcanic eruption. The "volcano" (a plastic 35 mm film cannister) erupts (the lid blows off) when gas pressure generated by dissolving alka seltzer is sufficiently high. It is realistic in that the timing of the eruption is difficult to predict precisely and in that the eruption occurs when the pressure of the gas exceeds the confining pressure of the lid. The experiment can be modified to show that an eruption will not occur if there is not enough gas pressure generated (small piece of alka seltzer) or if gas is allowed to escape gradually through holes punched in lid of film cannister.

Background

Not all volcanoes erupt explosively and unpredictably. The eruptive style (quiet streams of lava versus violent blasts of gases, ash, and debris) and eruption frequency and predictability are related to the viscosity (its resistance to flow) and amount of dissolved gas in the magma. (Magma = lava prior to eruption). Hot, runny magmas with little dissolved gas tend to flow smoothly out of vents and produce a volcano that is broad and not steep, such as the Hawaiian volcanoes.

On the other hand, slightly cooler magma with a higher dissolved gas content is much more viscous. Instead of running out smoothly, the magma may ooze out like toothpaste, clogging the vent. These volcanoes are steeper and have the typical "volcano shape" of famous volcanoes such as Mt. Fuji in Japan. Under this type of volcano, gases dissolved in magma can separate as

the pressure decreases as the magma rises closer to the Earth's surface. If the gases separate rapidly and cannot escape immediately, they can build up pressure greater than that of the overlying rock. When this happens, they break the rock suddenly as a violent explosion occurs sending a plume of gas and ash upwards to heights as great as 20 miles. Commonly there is little if any lava extruded. The ejecta may consist primarily of ash, pumice (instantaneously cooled magma containing abundant trapped air), and debris blown off the volcano by the eruption. The force of the eruption can blast material tens of miles from the volcano, causing extensive loss of life and damage. It may take thousands of years for sufficient pressure to build to cause an eruption, and it is difficult to predict when a volcano that has long been dormant will become active. If the pressure were released gradually it would not build up to the point where it could cause an explosive eruption. Because they erupt infrequently, unpredictably, and violently, and because they occur in populated areas (e.g. Japan, Indonesia, Philippines, Pacific northwest of the United States, Central and South America), these explosive volcanoes pose the greatest danger to humans.

Materials

- Plastic film cannisters with lids (black cylinders with gray lids are best) typically free from photo processing services
- Water
- Graduated cylinders (small)
- Alka seltzer (regular strength) tablets, cut into halves and quarters
- Lots of paper towels for clean up

Procedures

1. Put about 20 ml of water into cannister (Don't have to be exact.)
2. Add quarter tablet of alka seltzer and quickly put on cap.
3. Observe what happens for about 2 minutes.
4. Repeat using half of an alka seltzer tablet. (Watch out!)
5. Repeat using half of a tablet and lid with perforations that allow gas to escape.

Results and Discussion

1. *What happens?* The lid on cannister containing a quarter tablet will balloon upwards as the pressure increases, then deflate as it decreases, or it may possibly pop off gently. The lid on the cannister containing a half of a tablet will bulge and then blow off surprisingly violently, rising several feet in the air or travelling laterally 5 or 6 feet. The perforated lids will not blow off.
2. *Why?* There is not enough gas generated by one quarter of a tablet of alka seltzer to build sufficient pressure to blow the lid off. However, the gas generated by a half of an alka seltzer tablet is plenty to blow the lid part way across the room, as long as the lid is put on quickly and firmly so that no gas escapes. Students can use the ballooning of the lid prior to its blowing off to predict when the lid will blow off. This is analogous to volcanologists predicting when an eruption will occur based on the measurable bulging of a volcano. If the gas is allowed to escape gradually through holes in the lid, the gas

pressure from the dissolving alka seltzer will not build up to the point where it exceeds the confining pressure of the lid, and the lid will not blow off.

Additional Activities

Repeat each experiment several times. Using a second hand on a clock, time how long it takes the lid to blow off with the half tablet. Measure the distance that the lid goes. Data from several groups can be pooled, tabulated, and histograms made.

Older students could research the eruptive history of a particular volcano, the effects it has had on humankind, and its potential for future destruction. Some famous volcanoes: Kilauea and Mauna Loa (both "nonviolent" volcanoes on the island of Hawaii), Mt. Vesuvius, Mt. Pinatubo (erupted 1991), Mt. Rainier, Mt. Lassen, Mt. St. Helens, Tambora, Krakatoa, El Chichon, Nevado del Ruiz (erupted 1985 causing 23,000 deaths in Columbia). Locate these volcanoes on a world map (geography) and have students discover some dramatic story or event for their assigned volcano.

Watch "Volcano!", 1988, National Geographic video (58 min.). It is sobering for students to learn that the volcanologists featured on the video, the Kraffts, a French husband and wife team, were killed in the 1991 eruption of Unzen in Japan.